



Effects of whole body vibration in patients with chronic obstructive pulmonary disease – A randomized controlled trial[☆]

Rainer Gloeckl^{a,*}, Inga Heinzelmann^a, Sandra Baeuerle^b, Eva Damm^c, Anna-Lena Schwedhelm^c, Merve Diril^c, David Buhrow^c, Andreas Jerrentrup^c, Klaus Kenn^a

^a Department of Respiratory Medicine, Schoen Klinik Hospital, Schoenau am Koenigssee, Germany

^b Department of Sports Science, University of Mainz, Mainz, Germany

^c Department of Internal Medicine and Respiratory Medicine, University Marburg, Marburg, Germany

Received 12 September 2011; accepted 31 October 2011

Available online 21 November 2011

KEYWORDS

Chronic obstructive pulmonary disease;
Pulmonary rehabilitation;
Exercise;
Training;
Whole body vibration

Summary

Introduction: To date endurance and strength training are established and evidence-based exercise methods in patients with chronic obstructive pulmonary disease (COPD). There is an unmet need for further research in new and complementary exercise modalities. Additional whole body vibration training during pulmonary rehabilitation may be such a new approach that has not yet been investigated in patients with COPD.

Methods: Eighty-two patients (65 ± 9 yrs, FEV₁ pred. 38 ± 11%, female 51%) with COPD in GOLD stage III to IV assessed for a 3-week inpatient multidisciplinary rehabilitation program were on top randomly assigned to one of two intervention groups: (1) 3 × 3 min of bilateral dynamic squat exercises on a side-alternating vibration platform at 24–26 Hz three times per week (WBV) and (2) a control group (CON) with the same amount of exercise time without WBV.

Results: Thirty-six patients completed the study in each group. The improvement in 6-min walking distance was significantly higher in the WBV-group when compared to the CON-group (WBV: 64 ± 59 m, CON: 37 ± 52 m with a between-group difference of 27 m [95% CI, 1–53], *p* = 0.046). The time required for a sit-to-stand test also decreased more markedly in the WBV-group than in the CON-group (WBV: −4.0 ± 4.8 s, CON: −2.0 ± 3.1 s with

Abbreviations: 6-MWD, Six-minute-walking-distance; ATS, American Thoracic Society; BODE, The Body-Mass Index Airflow Obstruction Dyspnea and Exercise Capacity Index; BMI, Body-Mass-Index; CON, Control-group; COPD, Chronic obstructive pulmonary disease; CRQ, Chronic Respiratory Questionnaire; DLCO, Diffusion capacity of the lung for carbon monoxid; FEV₁, Forced expiratory volume in 1 s; GOLD, Global Initiative for Chronic Obstructive Lung Disease; HRQL, Health related quality of life; IQR, Interquartile range; LTOT, Long term oxygen therapy; MCID, Minimal clinical important difference; SST, Sit-to-stand test; WBV, Whole body vibration.

[☆] This trial has been registered in the Clinical Trials registry (identification number NCT01380639).

* Corresponding author. Tel.: +49 8652931871.

E-mail address: rainer.gloeckl@gmx.de (R. Gloeckl).

a between-group difference of -1.9 s [95% CI, -4.0 to 0.1], $p = 0.067$). Improvements in health-related quality of life were similar in both groups.

Conclusions: WBV training seems to be a promising new exercise modality for patients with COPD and may enhance the effects of a multidisciplinary rehabilitation program.

© 2011 Elsevier Ltd. All rights reserved.

Introduction

Limited exercise capacity is one of the main systemic manifestations of chronic obstructive pulmonary disease (COPD) associated with poor health-related quality of life (HRQL), exacerbations and increased mortality risk.^{1–3} Pulmonary rehabilitation is regarded as an evidence-based treatment for symptomatic patients with COPD to improve exercise tolerance, reduce symptoms of dyspnea and increase HRQL.^{4,5} Exercise training is seen as a cornerstone in pulmonary rehabilitation programs.⁶ Whilst the use of endurance and strength training are established and evidence-based exercise modalities in patients with COPD,^{7,8} the validation of further complementary exercise methods is considered a remaining challenge.⁹

Exercising on a whole body vibration plate might be such a new and additional approach. During a whole body vibration (WBV) session, subjects exercise on a vibrating platform that produces sinusoidal oscillations. In healthy subjects it has been shown that performing resistance training on a vibrating platform induces greater neuromuscular and hormonal responses than resistance training alone by providing a more intense stimulus.¹⁰ So far WBV has also been established in rehabilitation programs for patients with neurological diseases to improve postural control or for patients with osteoporosis to enhance bone mineral density.^{11–13} Up to now the effects of WBV training in patients with pulmonary diseases have rarely been studied. Based on the assumption that an increase in muscle strength and performance are considered the major effects of WBV,^{14,15} this could be a promising exercise modality for patients with COPD. Therefore we conducted this prospective randomized controlled trial to investigate the effects of WBV during multidisciplinary inpatient rehabilitation in patients with COPD.

Methods and materials

Study design

Consecutive patients admitted to a 3-week inpatient rehabilitation program were asked to participate in this randomized controlled parallel group study. Patients were not blinded to the intervention as this was not possible within this setting. Inclusion ran from April 2010 until April 2011. The trial was submitted to the Clinical Trials registry (identification number NCT01380639) and was approved by the Bavarian ethics committee (identification number 11050).

Study population

Eighty-two patients participated in this study. To attend the study, patients had to meet the following inclusion criteria:

(1) assured diagnosis of COPD stage III or IV according to the Global Initiative for Chronic Obstructive Lung Disease (GOLD)¹⁶ and (2) given written informed consent. Exclusion criteria were defined as follows: 1) status post surgery or post bone fracture, 2) status post deep vein thrombosis, 3) existing arterial aneurysm, 4) COPD exacerbation within the last 4 weeks or 5) non compliance to study protocol.

Intervention

Patients followed a multidisciplinary inpatient rehabilitation program of 3-weeks duration that included therapy units on 5 days per week. The rehabilitation program consisted of medical care, breathing therapy, education, nutritional counseling and psychological support. Exercise training consisted of endurance training (15 min cycling at 60% peak Watt) and strength training (4–6 exercises with 3 sets at the 20-repetition-maximum for major muscle groups). These components were equal for all patients. Additionally patients were randomly assigned to one of two intervention groups. Both groups attended a supervised squat exercise program. The only difference between groups was the surface on which the squats were performed. The first group (WBV), $n = 42$ performed squat exercises on a whole body vibration plate (GALILEO[®], Novotec Medical GmbH, Pforzheim, Germany) at 24–26 Hz and 6 mm peak-to-peak amplitude (Fig. 2). The side-alternating movement of the GALILEO[®] plate evokes muscle contractions on the entire flexor and extensor chain of muscles in the legs and all the way up to the trunk.^{17,18} Instead of voluntary muscle control like in common resistance training, the muscle contractions during vibration training (above 12 Hz) are caused by stretch reflexes.^{19,20} The user has no direct influence on muscle activity itself and can only control body posture, movement and exercise objective.

The second group, regarded as control group (CON), $n = 40$ performed the same amount of supervised squat training on the floor. The training schedule for both groups consisted of three sets with 3-min sessions of self-paced squats 3 times per week.

Outcomes and measurements

Primary outcome parameter of the study was the change in 6-min-walking-distance (6-MWD). The 6-MWD was performed according to the guidelines of the American Thoracic Society (ATS).²¹ The best out of two tests on day 1 and 2 was used as a baseline value to exclude any learning effects.²²

As a second exercise test a sit-to-stand test was performed. For the sit-to-stand test subjects were asked to stand up and sit down from a standard chair (height 43 cm) as quickly as possible with their arms folded across their chest. The time that patients needed to perform 5




Figure 1 Study flow diagram.

repetitions in a row was measured.²³ As the sit-to-stand test examines a body movement which is very closely connected to activities of daily living it is regarded as an appropriate tool to determine functional status in patients with COPD.²⁴

Lung function was measured with a Master Screen Bodyplethysmograph (Jaeger, Wuertzburg, Germany) in accordance with the ATS guidelines.^{16,25}

General HRQL was assessed by self-administered German-language version of the Chronic Respiratory Questionnaire (CRQ).²⁶ The CRQ is a widely used instrument in pulmonary rehabilitation and measures the dimensions of dyspnea, fatigue, emotional function and the ability to cope with COPD (mastery). The scores in each domain range from 1 (most severe impairment) to 7 (no impairment).

The BODE-index, a simple and established multidimensional grading system was used to assess the degree of disease related impairment and to predict the risk of mortality among patients with COPD.² The BODE-index is made up of a 10-point scale in which higher scores indicate an advanced stage of COPD and a higher risk of death. It consists of the Body-Mass-Index (B), the degree of airflow obstruction [FEV₁] (O), the degree of dyspnea [MMRC scale] (D) and exercise capacity (E) measured by the 6-MWD.

All measurements were performed on day 1 and 21 of the rehabilitation program.

Statistical analysis

Data are presented as mean and standard deviation or mean and 95%-confidence interval unless otherwise stated. Within-group differences were analyzed using paired *t* test. Between-group comparisons of changes in outcome parameters were carried out by using an unpaired *t* test. For unevenly distributed data the Wilcoxon's signed rank test (intra-group) and the Mann-Whitney-*U*-test (inter-group) were applied. The level of significance was set at $p < 0.05$. All data analyzes were performed using PASW Statistics Version 18.0 (Chicago, IL, USA).

We determined that a sample size of 36 patients in each group would show the inferiority of WBV-training, assuming inferiority at the margin of the minimal clinically important difference (MCID) of 35 m²⁷ with an estimated standard deviation of 53 m in the 6-MWD between groups to yield a power of 80% at a one-sided alpha level of 2.5%.

Randomization procedure was performed by using two permuted blocks of equal length with a ratio of 1:1.

We handled missing data as follows. CRQ-data for 1 patient in each group were missing. These missing values were excluded from the final analysis. There were no missing data for exercise capacity, lung function or the BODE-index.



Figure 2 Squat exercise on the whole body vibration platform GALILEO®.

Results

Due to 10 out of 82 patients dropped out of the study (for reasons see Fig. 1) 72 patients completed the study and were considered for the final per-protocol analysis. Thirty-six out of 42 patients in the WBV-group (86%) and 36 out of 40 in the CON-group (90%). Randomization strategy allocated patients into two comparable groups (for baseline values see Table 1). Pulmonary rehabilitation was effective in both groups as assessed by a significant increase in 6-MWD of 64 ± 59 m ($p < 0.01$) in the WBV-group as well as 37 ± 52 m ($p < 0.01$) in the CON-group with a between-group difference of 27 m [95% CI, 1–53], $p = 0.046$ (Fig. 3). Patients in both groups increased their 6-MWD above the MCID of 35 m.²⁷ Patients also improved their exercise capacity in the sit-to-stand test as shown by a decreased test time (WBV-group: -4.0 ± 4.8 s, CON-group: -2.0 ± 3.1 s, between-group difference of -1.9 s [95% CI, -4.0 to 0.1], $p = 0.067$) (Fig. 3). After pulmonary rehabilitation HRQL improved significantly within both groups. Almost all items of the CRQ (except for *mastery* in the CON-group) improved more than the MCID of 0.5 points²⁸ (CRQ total score in the WBV-group: 3.09 ± 4.28 points and in the CON-group: 2.92 ± 3.13 points, between-

group difference of 0.16 points [95% CI, -1.63 to 1.95], $p = 0.86$). For detailed changes in the CRQ scores see Fig. 4. The BODE-index decreased significantly and to a similar proportion in both groups (WBV-group: -0.7 ± 1.3 and in the CON-group: -0.9 ± 1.8 , between-group difference: 0.2 [95% CI, -0.8 to 0.8], $p = 0.91$) (Table 2).

No serious adverse events such as injuries, cardiac events or an increase in respiratory symptoms were observed according to the study protocol.

Discussion

The idea of vibration training came up in the early 1970s in order to train Russian cosmonauts and prevent loss of bone mineral and muscle mass during space flights. In the 1990s vibration training was rediscovered for exercise training in professional athletes. During the past decade there has been increased interest in WBV for therapeutic uses. So far several meta-analysis and systematic reviews have investigated the effects of WBV with inconsistent results.^{29,30} To date many studies concerning WBV research are methodologically weak and should be interpreted with caution.³¹ Study protocols have used widely variable WBV parameters and most included only small sample sizes which complicate interpretation of these studies. Some but not all of the studies reported similar improvements in muscle performance, balance, and bone mineral density for WBV when compared to traditional exercise programs.^{32–37}

Thus far only two studies have investigated the effects of WBV in patients with pulmonary diseases. In 2008 a German workgroup studied 8 patients with cystic fibrosis during a home-based training program on a WBV platform without a control group. After 6 months of training only a small increase in leg muscle power of 4.7% (range -16.4 to 74.5) could be found.³⁸ The second study investigated WBV versus conventional resistance training in 59 patients with COPD (FEV₁ pred. $47 \pm 21\%$).³⁹ WBV was found to be a promising training modality, which yielded improvements of similar magnitude for exercise capacity, muscle force and quality of life as those obtained with conventional resistance training after 12 weeks of pulmonary rehabilitation.

Therefore we conducted this study to compare two exercise protocols implemented within an inpatient rehabilitation program of 3-weeks duration in patients with more advanced COPD. The aim of this investigation was to assess if supplemental exercise training on a vibration platform leads to superior effects when compared to a conventional multidisciplinary rehabilitation program. As most patients with COPD suffer from a decreased muscle mass and power,⁴⁰ WBV training could be a promising new exercise modality.

The mean improvement in exercise capacity (6-MWD) is comparable to the magnitude that was reported in former pulmonary rehabilitation programs of comparable durations.^{41,42} However it is striking that patients in the WBV-group were able to improve their functional capacity to a significantly and considerably greater extent when compared to the CON-group. This is specifically remarkable as patients in both groups also performed supplemental endurance and strength training. The outcome of the sit-to-stand

Table 1 Baseline characteristics.

Characteristics	Patients randomized to rehabilitation		Patients completing rehabilitation	
	WBV-group (n = 42)	CON-group (n = 40)	WBV-group (n = 36)	CON-group (n = 36)
Age, yrs	64 ± 11	65 ± 7	64 ± 11	65 ± 7
Gender, female	23(55)	19(48)	18(50)	17(47)
LTOT, n	14(33)	15(38)	11(31)	13(36)
BMI, kg/m ²	25 ± 5	26 ± 6	24 ± 5	27 ± 6
BODE-index	4.3 ± 2.1	4.0 ± 2.0	4.0 ± 1.8	4.0 ± 2.0
6-MWD, m	342 ± 121	345 ± 105	349 ± 112	344 ± 104
6-MWD, % pred.	55 ± 19	57 ± 17	56 ± 18	57 ± 16
SST, sec.	15.7 ± 7.5	16.0 ± 7.5	15.7 ± 7.8	16.0 ± 7.5
FEV ₁ , % pred.	39 ± 11	38 ± 12	39 ± 11	38 ± 12
D _{LCO}	41 ± 16	43 ± 20	41 ± 17	44 ± 21
CRQ				
Total score	15.57 ± 3.29	14.34 ± 3.78	15.57 ± 3.29	14.57 ± 3.65
Dyspnea	3.42 ± 1.03	3.27 ± 1.33	3.42 ± 1.03	3.37 ± 1.30
Fatigue	4.03 ± 0.92	3.60 ± 1.02	4.03 ± 0.92	3.64 ± 1.02
Emotional function	4.20 ± 1.20	3.72 ± 1.20	4.21 ± 1.22	3.76 ± 1.21
Mastery	4.27 ± 1.25	4.11 ± 1.32	4.27 ± 1.25	4.23 ± 1.26

Data are presented in mean ± SD or number (%); WBV = whole body vibration, CON = control, LTOT = long-term oxygen therapy, BMI = body-mass-index, BODE-index = BMI-obstruction-dyspnea-exercise capacity-index; 6-MWD = 6-min-walking-distance, SST = sit-to-stand test, FEV₁ = forced expiratory volume (1 s), CRQ = chronic respiratory questionnaire.

test provided similar results in favor of the WBV-group although the between-group difference just failed to reach significance. We assume that these improvements in exercise capacity after WBV may be related to an increase in neuromuscular activation. One possibility for the WBV effect is through the vibratory stretch reflex in which the mechanical vibration elicits a myostatic stretch reflex mediated by the muscle spindle and Ia-afferents.⁴³ Other possible mechanisms of WBV benefits include enhancing postural control and improving quality of intermuscular coordination like the complex interplay of agonists and antagonists which are often disabled in patients with COPD.^{44–46}

Despite of patients' better exercise capacity in the WBV-group there was no transfer to an additional improvement in HRQL or the BODE-index. The reason for this could be that at the beginning of the study patients in the WBV-group already had a remarkable higher total score in the CRQ (+0.94 points) as well as a higher score at the BODE-index (+0.4 points) than patients in the CON-group. We guess that this difference at baseline may have yielded a potential underestimation of the WBV-group results regarding these two secondary outcome parameters. As patients still perceive the burden of dyspnea after pulmonary rehabilitation it remains debatable whether the objective measured superior improvements in exercise capacity in the WBV-group may also be reflected by the subjective measurement method of a questionnaire.

This randomized controlled trial is also associated with some limitations. Although all exercise sessions were supervised and advices for a high standard of squat performance were given, we did not record the number of repetitions patients performed during their squat training. This may have yielded potential differences between individuals. Nevertheless we chose the self-pace approach to

increase exercise feasibility. Another shortcoming may be that patients were not enrolled in a follow-up investigation, as our study did not focus on the long-term effects. Also the unblinded nature of the study protocol may have yielded a certain bias due to a placebo effect. Nevertheless it was not possible to blind patients or researchers to the exercise intervention.

In consideration of the short total exposure time to WBV (81 min in 3 weeks), the additionally attained effects are quite striking. Nevertheless, WBV training should not be considered as an alternative exercise modality which can be used instead of "normal" exercise training. Specific exercises for arms, accessory respiratory muscles or the upper trunk cannot be easily performed on a vibration plate and it is not possible to undertake endurance training on a vibration platform.

In conclusion, WBV training seems to be an effective and feasible exercise modality for patients with advanced COPD that may even enhance the effects of a comprehensive multidisciplinary rehabilitation program. Further studies are needed in order to define optimal intensity and duration of WBV as well as to investigate its possible long-term effects.

Author contributions

RG: contributed to the organization of the study, study design, data analysis, writing the manuscript and sharing scientific discussions.

IH: contributed to the organization of the study, study design and sharing scientific discussions.

SB: contributed to the organization of the study and data collection.




Figure 3 Comparison of exercise capacity (boxes with 95% confidence intervals represent point estimate for the differences between groups) after a 3-week multidisciplinary rehabilitation program with an additional squat exercise program on the floor (control group) or a squat exercise program during whole body vibration (WBV-group). Patients in the WBV-group increased their 6-minute-Walking distance by 27 m superior to the control group. Patients in the WBV-group decreased their time needed to perform the five repetition sit-to-stand test by 1.9 s superior to the patients in the control group.




Figure 4 Comparison of health-related quality of life gathered by chronic respiratory questionnaire [CRQ] (boxes with 95% confidence intervals represent point estimate for the differences between groups) after a 3-week multidisciplinary rehabilitation program with an additional squat exercise program on the floor (control group) or a squat exercise program during whole body vibration (WBV-group).

Table 2 Comparison of treatment effects.

Outcome	Per protocol			p value	Difference in intention-to-treat analysis (95% CI)
	Mean changes from baseline (SD)		Difference (95% CI)		
	WBV-group (n = 36)	CON-group (n = 36)			
Primary					
6-MWD, m	64.0 ± 59.1 ^a	37.3 ± 52.2 ^a	26.7 (0.5 to 52.9)	0.046	21.3 (−3.1 to 45.5)
6-MWD, % pred.	10.5 ± 10.2 ^a	6.1 ± 9.0 ^a	4.3 (−0.2 to 8.8)	0.061	3.4 (−0.7 to 7.6)
Secondary					
SST, sec.	−4.0 ± 4.8 ^{b,a}	−2.0 ± 3.1 ^a	−1.9 (−4.0 to 0.1)	0.067	−1.6 (−3.4 to 0.3)
FEV ₁ , % pred.	0.8 ± 6.6	2.6 ± 5.7 ^a	−1.8 (−4.7 to 1.1)	0.22	−1.7 (−4.2 to 0.9)
BODE-index	−0.7 ± 1.3 ^a	−0.9 ± 1.8 ^a	0.2 (−0.8 to 0.8)	0.91 ^c	0.1 (−0.7 to 0.7)
CRQ					
Total score	3.09 ± 4.28 ^a	2.92 ± 3.13 ^a	0.16 (−1.63 to 1.95)	0.86	0.10 (−1.61 to 1.63)
Dyspnea	0.67 ± 1.13 ^a	0.50 ± 0.84 ^a	0.18 (−0.32 to 0.68)	0.48	0.13 (−0.13 to 0.58)
Fatigue	0.69 ± 0.86 ^a	0.72 ± 0.87 ^a	−0.03 (−0.46 to 0.40)	0.88	−0.05 (−0.44 to 0.34)
Emotional function	0.70 ± 1.30 ^a	0.82 ± 0.68 ^a	−0.13 (−0.66 to 0.41)	0.64	−0.13 (−0.61 to 0.34)
Mastery	0.58 ± 1.20 ^a	0.33 ± 0.85 ^a	0.25 (−0.28 to 0.78)	0.34	0.20 (−0.26 to 0.66)

Data are presented in mean ± SD unless otherwise indicated; WBV = whole body vibration, CON = control, 6-MWD = 6-min-walking-distance, SST = sit-to-stand test; FEV₁ = forced expiratory volume (1 s), BODE-index = Bodymass-obstruction-dyspnea-exercise capacity-index; CRQ = chronic respiratory questionnaire.

^a Intra-group change of $p < 0.05$.

^b Wilcoxon Test.

^c Mann–Whitney U Test.

ED: contributed to the organization of the study and data collection.

AS: contributed to the organization of the study and data collection.

MD: contributed to the organization of the study and data collection.

DB: contributed to the organization of the study and data collection.

AJ: contributed to the study design and shared scientific discussions.

KK: contributed to the study hypothesis, study design, writing the manuscript and sharing scientific discussions.

Financial/nonfinancial disclosures

The authors have reported to *RESPIRATORY MEDICINE* that no potential conflicts of interest exist with any companies/organizations whose products or services may be discussed in this article.

Acknowledgment

The authors thank Mrs. Wanda Tamerl and her staff from the Department of Clinical Diagnostics for their assistance with data collection.

Conflict of interest

All authors (Rainer Gloeckl, Inga Heinzelmann, Sandra Baeuerle, Eva Damm, Anna-Lena Schwedhelm, Merve Diril,

David Buhrow, Andreas Jerrentrup, and Klaus Kenn) acknowledge that there are no conflicts of interest with any companies/organizations whose products or services may have influenced this study or manuscript.

Ethics statement

This trial was approved by the Bavarian ethics committee (identification number 11050).

References

- Swallow EB, Reyes D, Hopkinson NS, Man WD, Porcher R, Cetti EJ, et al. Quadriceps strength predicts mortality in patients with moderate to severe chronic obstructive pulmonary disease. *Thorax* 2007;**62**(2):115–20.
- Celli BR, Cote CG, Marin JM, Casanova C, Montes de Oca M, Mendez RA, et al. The body-mass index, airflow obstruction, dyspnea, and exercise capacity index in chronic obstructive pulmonary disease. *N Engl J Med* 2004;**350**(10):1005–12.
- Garcia-Aymerich J, Lange P, Benet M, Schnohr P, Anto JM. Regular physical activity reduces hospital admission and mortality in chronic obstructive pulmonary disease: a population based cohort study. *Thorax* 2006;**61**(9):772–8.
- Lacasse Y, Goldstein R, Lasserson TJ, Martin S. Pulmonary rehabilitation for chronic obstructive pulmonary disease. *Cochrane Database Syst Rev* 2006;(4):CD003793.
- Kenn K, Sczepanski B. Pulmonary rehabilitation before and after lung transplantation. *Pneumologie* 2011;**65**(7):419–27.
- Ries AL, Bauldoff GS, Carlin BW, Casaburi R, Emery CF, Mahler DA, et al. Pulmonary rehabilitation: joint ACCP/AACVPR evidence-based clinical practice guidelines. *Chest* 2007;**131**(5 Suppl):45–42S.

7. Langer D, Hendriks E, Burtin C, Probst V, van der Schans C, Paterson W, et al. A clinical practice guideline for physiotherapists treating patients with chronic obstructive pulmonary disease based on a systematic review of available evidence. *Clin Rehabil* 2009;23(5):445–62.
8. Nici L, ZuWallack R, Wouters E, Donner CF. The ATS/ERS statement on pulmonary rehabilitation. *Eur Respir J* 2006;28(3):461–2.
9. Troosters T, Gosselink R, Janssens W, Decramer M. Exercise training and pulmonary rehabilitation: new insights and remaining challenges. *Eur Respir Rev* 2010;19(115):24–9.
10. Ronnestad BR. Comparing the performance-enhancing effects of squats on a vibration platform with conventional squats in recreationally resistance-trained men. *J Strength Cond Res* 2004;18(4):839–45.
11. Zha DS, Zhu QA, Pei WW, Zheng JC, Wu SH, Xu ZX, et al. Does whole-body vibration with alternative tilting increase bone mineral density and change bone metabolism in senior people? *Aging Clin Exp Res* 2011 [Epub ahead of print].
12. Von Stengel S, Kemmler W, Bebenek M, Engelke K, Kalender WA. Effects of whole-body vibration training on different devices on bone mineral density. *Med Sci Sports Exerc* 2011;43(6):1071–9.
13. Merkert J, Butz S, Nieczaj R, Steinhagen-Thiessen E, Eckardt R. Combined whole body vibration and balance training using vibrosphere(R): improvement of trunk stability, muscle tone, and postural control in stroke patients during early geriatric rehabilitation. *Z Gerontol Geriatr* 2011 [Epub ahead of print].
14. Petit PD, Pensini M, Tessaro J, Desnuelle C, Legros P, Colson SS. Optimal whole-body vibration settings for muscle strength and power enhancement in human knee extensors. *J Electromyogr Kinesiol* 2010;20(6):1186–95.
15. Rittweger J. Vibration as an exercise modality: how it may work, and what its potential might be. *Eur J Appl Physiol* 2010;108(5):877–904.
16. Rabe KF, Hurd S, Anzueto A, Barnes PJ, Buist SA, Calverley P, et al. Global strategy for the diagnosis, management, and prevention of chronic obstructive pulmonary disease: GOLD executive summary. *Am J Respir Crit Care Med* 2007;176(6):532–55.
17. Torvinen S, Kannu P, Sievanen H, Jarvinen TA, Pasanen M, Kontulainen S, et al. Effect of a vibration exposure on muscular performance and body balance. Randomized cross-over study. *Clin Physiol Funct Imaging* 2002;22(2):145–52.
18. Wirth B, Zurfluh S, Muller R. Acute effects of whole-body vibration on trunk muscles in young healthy adults. *J Electromyogr Kinesiol* 2011;21(3):450–7.
19. Cardinale M, Rittweger J. Vibration exercise makes your muscles and bones stronger: fact or fiction? *J Br Menopause Soc* 2006;12(1):12–8.
20. Cardinale M, Lim J. Electromyography activity of vastus lateralis muscle during whole-body vibrations of different frequencies. *J Strength Cond Res* 2003;17(3):621–4.
21. ATS-statement. Guidelines for the six-minute walk test. *Am J Respir Crit Care Med* 2002;166(1):111–7.
22. Hernandez NA, Wouters EF, Meijer K, Annegarn J, Pitta F, Spruit MA. Reproducibility of 6-minute walking test in patients with COPD. *Eur Respir J* 2010.
23. Bohannon RW. Reference values for the five-repetition sit-to-stand test: a descriptive meta-analysis of data from elders. *Percept Mot Skills* 2006;103(1):215–22.
24. Canuto FF, Rocco CC, de Andrade DV, Sampaio LM, Oliveira CS, Correa FI, et al. Neurophysiological comparison between the sit-to-stand test with the 6-minute walk test in individuals with COPD. *Electromyogr Clin Neurophysiol* 2010;50(1):47–53.
25. Macintyre N, Crapo RO, Viegi G, Johnson DC, van der Grinten CP, Brusasco V, et al. Standardisation of the single-breath determination of carbon monoxide uptake in the lung. *Eur Respir J* 2005;26(4):720–35.
26. Puhan MA, Behnke M, Frey M, Grueter T, Brandli O, Lichtenschopf A, et al. Self-administration and interviewer-administration of the German chronic respiratory questionnaire: instrument development and assessment of validity and reliability in two randomised studies. *Health Qual Life Outcomes* 2004;2:1.
27. Puhan MA, Mador MJ, Held U, Goldstein R, Guyatt GH, Schunemann HJ. Interpretation of treatment changes in 6-minute walk distance in patients with COPD. *Eur Respir J* 2008;32(3):637–43.
28. Schunemann HJ, Puhan M, Goldstein R, Jaeschke R, Guyatt GH. Measurement properties and interpretability of the chronic respiratory disease questionnaire (CRQ). *Copd* 2005;2(1):81–9.
29. Conway GE, Szalma JL, Hancock PA. A quantitative meta-analytic examination of whole-body vibration effects on human performance. *Ergonomics* 2007;50(2):228–45.
30. Nordlund MM, Thorstenson A. Strength training effects of whole-body vibration? *Scand J Med Sci Sports* 2007;17(1):12–7.
31. Cardinale M, Wakeling J. Whole body vibration exercise: are vibrations good for you? *Br J Sports Med* 2005;39(9):585–9. discussion 9.
32. Mikhael M, Orr R, Fiatarone Singh MA. The effect of whole body vibration exposure on muscle or bone morphology and function in older adults: a systematic review of the literature. *Maturitas* 2010;66(2):150–7.
33. Merriman H, Jackson K. The effects of whole-body vibration training in aging adults: a systematic review. *J Geriatr Phys Ther* 2009;32(3):134–45.
34. Stamm C. Whole body vibration - a systematic review. *J Physiotherapie [GER]* 2010;62/9:6–12.
35. Tihanyi TK, Horvath M, Fazekas G, Hortobagyi T, Tihanyi J. One session of whole body vibration increases voluntary muscle strength transiently in patients with stroke. *Clin Rehabil* 2007;21(9):782–93.
36. Osawa Y, Oguma Y. Effects of resistance training with whole-body vibration on muscle fitness in untrained adults. *Scand J Med Sci Sports* 2011 [Epub ahead of print].
37. Verschueren SM, Roelants M, Delecluse C, Swinnen S, Vanderschueren D, Boonen S. Effect of 6-month whole body vibration training on hip density, muscle strength, and postural control in postmenopausal women: a randomized controlled pilot study. *J Bone Miner Res* 2004;19(3):352–9.
38. Rietschel E, van Koningsbruggen S, Fricke O, Semler O, Schoenau E. Whole body vibration: a new therapeutic approach to improve muscle function in cystic fibrosis? *Int J Rehabil Res* 2008;31(3):253–6.
39. Salhi B, van Meerbeeck J, Joos GF, Derom E. *Effects of whole body vibration in patients with COPD: a randomized study.* ATS conference 2011. Denver: A17192.
40. Seymour JM, Spruit MA, Hopkinson NS, Natanek SA, Man WD, Jackson A, et al. The prevalence of quadriceps weakness in COPD and the relationship with disease severity. *Eur Respir J* 2010;36(1):81–8.
41. von Leupoldt A, Hahn E, Taube K, Schubert-Heukeshoven S, Magnussen H, Dahme B. Effects of 3-week outpatient pulmonary rehabilitation on exercise capacity, dyspnea, and quality of life in COPD. *Lung* 2008;186(6):387–91.
42. Puhan MA, Busching G, Schunemann HJ, VanOort E, Zaugg C, Frey M. Interval versus continuous high-intensity exercise in chronic obstructive pulmonary disease: a randomized trial. *Ann Intern Med* 2006;145(11):816–25.
43. Nishihira Y, Iwasaki TAH. Effects of whole body vibration stimulus and voluntary contraction on motorneuron pool. *Adv Exerc Sports Physiol* 2002;8(4):83–6.
44. Bosco C, Colli R, Introini E, Cardinale M, Tsarpela O, Madella A, et al. Adaptive responses of human skeletal muscle to vibration exposure. *Clin Physiol* 1999;19(2):183–7.

45. Kawanabe K, Kawashima A, Sashimoto I, Takeda T, Sato Y, Iwamoto J. Effect of whole-body vibration exercise and muscle strengthening, balance, and walking exercises on walking ability in the elderly. *Keio J Med* 2007;56(1):28–33.
46. Butcher SJ, Meshke JM, Sheppard MS. Reductions in functional balance, coordination, and mobility measures among patients with stable chronic obstructive pulmonary disease. *J Cardiopulm Rehabil* 2004;24(4):274–80.